

Amendments to the Claims

Please enter the following amendments to the claims:

1. (Currently Amended) A method of determining a set of large sequences from an electronic data base comprising a set $D = \{d_1, \dots, d_n\}$ of transactions d_i ($1 \leq i \leq n$) in a computer system with an implemented query module, each of the large sequences on the set D of transactions d_i having a support value greater than or equal to a given support value S , each of the transactions d_i of the set D being a sequence of items of a record $E = \{e_1, \dots, e_m\}$ of items e_j ($1 \leq j \leq m$) and the method comprising the following steps:
 - a) determining a set L_1 of large sequences from the set D of transactions, the large sequences of set L_1 each comprising exactly one item of the record E , and an assigned support value S_{L_1} on the sequence D of transactions each being greater than or equal to the given support value S ;
 - b) determining a set L_2 of large sequences from the set D of transactions, the large sequences of set L_2 each comprising exactly two items of the record E in a respective order R_{L_2} , and an assigned support value S_{L_2} on the set D of transactions each being greater than or equal to the given support value S , and nothing but sequences comprising one of the large sequences of set L_1 , as a partial sequence, being taken into account in determining set L_2 ;
 - c) determining a set L_k ($k > 2$) of large sequences from the set D of transactions, the large sequences of set L_k each comprising exactly k items of record E in a respective order R_{L_k} , and an assigned support value S_{L_k} on the sequence D of transactions each being greater than or equal to the given support value S , and nothing but sequences comprising

- two of the large sequences of set L_{k-1} , as partly overlapping partial sequences, with the respective order $R_{L_{k-1}}$, being taken into account in determining set L_k ; and
- d) repeating step c) for $k = k+1$ and terminating the repetition of step c) when a given predefined termination condition is fulfilled.

2. (Original) The method as claimed in claim 1, wherein the set D of transactions is searched for candidate sequences which comprise two of the large sequences of set L_1 or set L_{k-1} , respectively, as partly overlapping partial sequences, in determining set L_2 in step b) and set L_k in step c), respectively, and wherein an assigned support value counter is registered for each candidate sequence found for the first time, and the assigned support value counter is incremented when the respective candidate sequence is determined again in searching the set D of transactions.

3. (Original) The method as claimed in claim 1, wherein:

- the set D of transactions is searched for candidate sequences comprising one of the large sequences of set L_1 , as a partial sequence, when determining set L_2 in step b), and an assigned support value counter is registered for each candidate sequence found for the first time, and the assigned support value counter is incremented when the respective candidate sequence is determined again in searching the set D of transactions; and
- a candidate set of sequences is formed, when determining set L_k ($k > 2$) in step c), which set comprises all combinations in pairs of the sequences of set L_{k-1} , and of the set D of transactions those transactions are defined which comprise at least

one sequence of the candidate set, and the support value is determined for those sequences of the candidate set for which a transaction was found that comprised this sequence.

4. (Previously amended) The method as claimed in claim 2, wherein a display value is generated and updated for each assigned support value counter to indicate the transaction for which the assigned support value counter was last incremented.
5. (Previously amended) The method as claimed in claim 3, wherein only those sequences of the candidate set of sequences corresponding to all combinations in pairs of the sequences of set L1 are taken into account of which the first item lies within a given range of values.
6. (Currently Amended) A computer program product for determining a set of large sequences from an electronic data base comprising a set $D = \{d_1, \dots, d_n\}$ of transactions d_i ($1 \leq i \leq n$) in a computer system with an implemented query module, each of the large sequences on the set D of transactions d_i having a support value greater than or equal to a given support value S, each of the transactions d_i of the set D being a sequence of items of a record $E = \{e_1, \dots, e_m\}$ of items e_j ($1 \leq j \leq m$) and the product comprising the following means:
 - a) means recorded on an electronic storage medium for determining a set L1 of large sequences from the set D of transactions, the large sequences of set L1 each comprising exactly one item of the record E, and an assigned support value S_{L1} on the sequence D of transactions each being greater than or equal to the given support value S;

- b) means recorded on the electronic storage medium for determining a set L_2 of large sequences from the set D of transactions, the large sequences of set L_2 each comprising exactly two items of the record E in a respective order R_{L_2} , and an assigned support value S_{L_2} on the set D of transactions each being greater than or equal to the given support value S , and nothing but sequences comprising one of the large sequences of set L_1 , as a partial sequence, being taken into account in determining set L_2 ;
- c) means recorded on the storage medium for determining a set L_k ($k > 2$) of large sequences from the set D of transactions, the large sequences of set L_k each comprising exactly k items of record E in a respective order R_{L_k} , and an assigned support value S_{L_k} on the sequence D of transactions each being greater than or equal to the given support value S , and nothing but sequences comprising two of the large sequences of set L_{k-1} , as partly overlapping partial sequences, with the respective order $R_{L_{k-1}}$, being taken into account in determining set L_k ; and
- d) means recorded on the electronic storage medium for repeating step c) for $k = k+1$ and terminating the repetition of step c) when a ~~given~~ predefined termination condition is fulfilled.

7. (Currently Amended) An integrated sequential analysis system, comprising:

- an electronic data base comprising a set $D = \{d_1, \dots, d_n\}$ of transactions d_i ($1 \leq i \leq n$), each of the large sequences on the set D of transactions d_i having a support value greater than or equal to a given support value S , each of the transactions d_i of the set D being a sequence of items of a record $E = \{e_1, \dots, e_m\}$ of items e_j ($1 \leq j \leq m$);
- a query module comprising a query means coupled to the data base and a processing

- means for detecting query parameters and generating queries to the query means;
- means for determining a set L_1 of large sequences from the set D of transactions, the large sequences of set L_1 each comprising exactly one item of the record E , and an assigned support value S_{L_1} on the sequence D of transactions each being greater than or equal to the given support value S ;
 - means for determining a set L_2 of large sequences from the set D of transactions, the large sequences of set L_2 each comprising exactly two items of the record E in a respective order R_{L_2} , and an assigned support value S_{L_2} on the set D of transactions each being greater than or equal to the given support value S , and nothing but sequences comprising one of the large sequences of set L_1 , as a partial sequence, being taken into account in determining set L_2 ;
 - means for determining a set L_k ($k > 2$) of large sequences from the set D of transactions, the large sequences of set L_k each comprising exactly k items of record E in a respective order R_{L_k} , and an assigned support value S_{L_k} on the sequence D of transactions each being greater than or equal to the given support value S , and nothing but sequences comprising two of the large sequences of set L_{k-1} , as partly overlapping partial sequences, with the respective order $R_{L_{k-1}}$, being taken into account in determining set L_k ; and
 - means for repeating step c) for $k = k+1$ and terminating the repetition of step c) when a given predefined termination condition is fulfilled.
8. (Previously Added) The method as claimed in claim 3, wherein a display value is generated and updated for each assigned support value counter to indicate the transaction for which the assigned support value counter was last incremented.

9. (Previously Added) The method as claimed in claim 4, wherein only those sequences of the candidate set of sequences corresponding to all combinations in pairs of the sequences of set L1 are taken into account of which the first item lies within a given range of values.
10. (Newly Added) The method as claimed in claim 1, where the predefined termination condition comprises substantially all large sequences of maximum length being found.
11. (Newly Added) The method as claimed in claim 1, where the predefined termination condition comprises reaching a maximum length of the large sequences.
12. (Newly Added) The method as claimed in claim 6, where the predefined termination condition comprises substantially all large sequences of maximum length being found.
13. (Newly Added) The method as claimed in claim 6, where the predefined termination condition comprises reaching a maximum length of the large sequences.
14. (Newly Added) The method as claimed in claim 7, where the predefined termination condition comprises substantially all large sequences of maximum length being found.
15. (Newly Added) The method as claimed in claim 7, where the predefined termination condition comprises reaching a maximum length of the large sequences.